



**PROVIDER'S REPORT AND TREATMENT ORDERS
FOR SEIZURE MANAGEMENT AT SCHOOL
CONFIDENTIAL**

Salem-Keizer School District 24J
Student Services
P.O. Box 12024
Salem, OR 97309-0024
(503) 399-3101

*PHYSICIAN'S AUTHORIZATION FOR ADMINISTRATION OF
SPECIALIZED PHYSICAL HEALTH CARE SERVICES*

PLEASE FAX TO STUDENT SERVICES AT (503) 316-3500

Student ID: _____ Student Name: _____ DOB: _____

School Name: _____ Nurse: _____ may be reached at (503) 399-3101

IF YOU SEE THIS:	DO THIS:
<input type="checkbox"/> Convulsive Generalized Tonic Clonic: You will see loss of consciousness. Stiffening of the body. Rhythmic jerking movements. Convulsive seizures may last 1-5 minutes. The child may have a warning (aura) before the seizure. Sleepiness and confusion may occur after the seizure.	<ol style="list-style-type: none"> 1. Time the seizure 2. Keep calm. Provide reassurance. 3. Protect head, keep airway clear, turn on side if possible. 4. Do not place anything in mouth. 5. Call 911 if student is injured or has difficulty breathing. 6. Call parent. 7. Stay with student until recovered from seizure. 8. Administer rescue treatments as marked below and call 911 if not already done.
<input type="checkbox"/> Focal: These seizures may begin with an aura. They may be partly alert or unconscious. You may see lip smacking, chewing, eye blinking, or picking at clothes. These seizures usually last 1-2 minutes.	<ol style="list-style-type: none"> 1. Time the seizure 2. Gently guide child away from danger. 3. Stay with student and reassure them until recovered from seizure. 4. Do not treat staring that is stopped by a touch or a nudge. 5. Call parent. 6. Administer rescue treatments as marked below and call 911 if not already done.
<input type="checkbox"/> Absence: You will see quick changes in alertness. May see eye flutter or small twitching. Usually, lasts less than 10 seconds.	

RESCUE TREATMENTS

Child has a VNS. Child/staff may swipe with aura. Staff may swipe at onset of seizure and every 60 seconds until seizure stops.

Give rescue medications below if seizure does not stop within _____ minutes.

If seizure lasts longer than ____ minutes, administer:

<input type="checkbox"/> Diastat ____ mg rectally	<input type="checkbox"/> Midazolam ____ mg in the nose	<input type="checkbox"/> Clonazepam ____ mg in the cheek	<input type="checkbox"/> Other _____
---	--	--	--------------------------------------

Multistep seizure rescue plan — Please see attached letter for details.

If cluster of _____ or more seizures in ____ minutes, administer:

<input type="checkbox"/> Diastat ____ mg rectally	<input type="checkbox"/> Midazolam ____ mg in the nose	<input type="checkbox"/> Clonazepam ____ mg in the cheek	<input type="checkbox"/> Other _____
---	--	--	--------------------------------------

Multistep seizure rescue plan — Please see attached letter for details.

If no emergency medication is at school and the child is experiencing seizures:
 Call family to bring medications to school or pick up child. **Call 911 if seizure lasts more than _____ minutes.**

Other: _____

CALL 911 For:

- Absence of breathing and/or pulse (Start CPR)
- After emergency medication is administered
- For continued pale or bluish skin/lips after seizure has stopped
- For continued noisy breathing after the seizure has stopped
- Other: _____